arXiv:1305.0541v1 [physics.plasm-ph] 2 May 2013

Quantum Hydrodynamics Approach to The Research of Quantum Effects and Vorticity Evolution in Spin Quantum Plasmas

Trukhanova M. Iv.[a]

(Dated: 02.05.2013)

In this paper we explicate a method of magneto quantum hydrodynamics (MQHD) for the study of the quantum evolution of a system of spinning fermions in an external electromagnetic field. The fundamental equations of microscopic quantum hydrodynamics (the momentum balance equation and the magnetic moment density equation) were derived from the many-particle microscopic Schrodinger equation with $Spin-Spin$ and $Coulomb$ modified Hamiltonian. Using the developed approach the vorticity evolution equation and quantum spin vorticity dynamics equation for the quantum spinning plasma were derived. The effects of new spin forces and spin-spin interaction contributions on the motion of fermions, evolution of the magnetic moment density and vorticity generation were predicted. This results can be used for the theoretical studies of spinning many-particle systems, especially dense quantum plasmas in compact astrophysical objects, plasmas in semiconductors and micro-mechanical systems, in quantum x-ray free-electron lasers.

PACS numbers: 52.35.We, 67.10.-j
Keywords: Quantum hydrodynamics, spin-spin Interactions, vorticity, spin quantum plasmas

The spinning quantum fluid plasma is becoming of increasing current interest[1] -[7]. Hydrodynamics equations of a spinning fluid for the Pauli equation with the quantum particle angular momentum spin was presented since the pioneering works by Takabayashi and Vigier[8] -[11]. The vector representation of non-relativistic spinning particle leads to appearance of new quantum effects had been separated as non-linear terms which arises from the inhomogeneity of spin distribution.

The quantum effects in plasma can be represented by three main quantum corrections. The first is a quantum force, the multiparticle quantum Bohm or Madelung potential, proportional to powers of $\hbar$ and produced by density fluctuations[12],[13]. The another one associated with the quantum particle angular momentum spin by the possible inhomogeneity of the external and spin magnetic fields. In the momentum balance equation this force appears through the magnetization energy[1]. And the latter force associated only with the spin magnetic moment of the particle[8].

The most interesting and defining features of a quantum spinning plasmas can be derived from the vorticity equation. It had been derived by[14] that the vorticity, constructed from spin field of a quantum spinning plasma, combines with the classical generalized vorticity to yield a new grand generalized vorticity that obeys the standard vortex dynamics.

We also can see that the astrophysics becomes a rapidly growing field of research. It is important that the consequences of turbulent plasma movement in the solar photosphere leads to the generation of vorticity while magnetic vortices are produced by the magnetic tension. For example, MHD simulations of magnetoconvection had been used to analyse the generation of small-scale vortex motions in the solar photosphere. It had been shown that two different types of photospheric vorticity, magnetic and non-magnetic, are generated in the domain using the vorticity equation, combined with the G-band radiative diagnostics[15]. The presence of vortex motions for the astrophysics had been developed in[16] -[17].

The extracting coherent vortices out of turbulent flows had been applied to simulations of resistive drift-wave turbulence in magnetized plasma[18]. The quasi-hydrodynamic and quasi-adiabatic regimes had been investigated.

The formation and dynamics of dark solitons and vortices in quantum electron plasmas had been studied in[19]. A pair of equations comprising the nonlinear Schrdinger and Poisson system of equations, which conserves the number of electrons as well as their momentum and energy had been used. It had been shown that the gradient free-energy contained in equilibrium spin vorticity can cause electromagnetic modes, in particular the light wave[20].

The collective electron angular momentum spin effects in spinning quantum plasmas can be investigated using insights from quantum kinetic theory or some effective theory. We propose a method of quantum hydrodynamics which allows to obtain a description of the collective effects in magnetized quantum plasmas in terms of functions in the physical space.

A quantum mechanics description for systems of N interacting spinning particles is based upon the many-particle Schrodinger equation (MPSE) that specifies a wave function in a 3N-dimensional configuration space. As wave processes, processes of information transfer and other spin transport processes occur in the three-dimensional physical space, a need arises to turn to a mathematical method of physically observable values which are determined in a 3D physical space. To do so

---

[a] M.V.Lomonosov Moscow State University, Faculty of Physics, Leninskie Gory, Moscow, Russia; Electronic mail: mar-tiv@yandex.ru

we should derive the fundamental equations those determine dynamics of functions of three variables, starting from MPSE. This problem has been solved with the creation of a method of many-particle quantum hydrodynamics (MPQHD).

In this article, using the MQHD approach we derive the fundamental balance equation, the magnetic moment evolution equation, the vorticity dynamics equation and the magnetic vorticity evolution equation for the magnetized quantum plasmas.

I. FUNDAMENTAL EQUATIONS OF THE FERMION QUANTUM HYDRODYNAMICS

In this section we derive the system of magneto quantum hydrodynamics (MQHD) equations for charged and neutral particles from the many-particle microscopic Schrodinger equation

$$i\hbar\frac{\partial\psi_s(R,t)}{\partial t}=(\hat{H}\psi)_s(R,t), \tag{1}$$

where $R=(\vec{r}_1,...,\vec{r}_N)$. We consider a system of N interacting fermions with equal masses m_j, charged and proper magnetic moments in an external electromagnetic field. A state of the system of N fermions is determined by a wave function in the 3N-dimensional configuration space, which is a $rank-N spinor$

$$\psi_s(R,t)=\psi_{s_1,s_2,....s_N}(\vec{r}_1,...,\vec{r}_N,t). \tag{2}$$

The Hamiltonian has the form

$$\hat{H}=\sum_{j=1}^{N}(\frac{\hat{D}_j^2}{2m_j}+q_j\varphi_{j,ext}-\mu_j\hat{\sigma}_j^\alpha B_{j,ext}^\alpha)+ \tag{3}$$

$$+\frac{1}{2}\sum_{j\neq k}^{N}q_jq_kG_{jk}-\frac{1}{2}\sum_{j\neq k,k}^{N}\mu_j^2F_{jk}^{\alpha\beta}\hat{\sigma}_j^\alpha\hat{\sigma}_k^\beta,$$

where $\mu_j=g\mu_B/2$, μ_{jB} - is the electron or positron magnetic moment and $\mu_{jB}=q_j\hbar/2m_jc$ - is the Bohm magneton, q_j stands for the charge of electrons $q_e=-e$ or for the charge of positrons $q_p=e$, and $\hbar$ -is the Planck constant, $g\simeq 2.0023193$. The covariant derivative operator is

$$\hat{D}_j^\alpha=-i\hbar\hat{\nabla}_j^\alpha-\frac{q_j}{c}A_j^\alpha, \tag{4}$$

where $\vec{A}_{ext},\varphi_{j,ext}$ - are the vector and scalar potentials of external electromagnetic field.

Green's functions of the *Coulomb* and $Spin-Spin$ interaction are

$$G_{ij}=\frac{1}{r_{jk}},\qquad F_{jk}^{\alpha\beta}=4\pi\delta_{\alpha\beta}\delta(\vec{r}_{jk})+\partial_j^\alpha\partial_k^\beta\frac{1}{r_{jk}}. \tag{5}$$

The first step in the construction of MQHD apparatus is to determine the concentration of particles in the neighborhood of $\vec{r}$ in a physical space. If we define the concentration of particles as quantum average of the concentration operator in the coordinate representation $\hat{\rho}=\sum_j\delta(\vec{r}-\vec{r}_j)$ we obtain

$$\rho(\vec{r},t)=\sum_S\int dR\sum_j^N\delta(\vec{r}-\vec{r}_j)\psi_s^+(R,t)\psi_s(R,t), \tag{6}$$

Differentiation of $\rho(\vec{r},t)$ with respect to time and applying of the Schrodinger equation with Hamiltonian 3 leads to continuity equation

$$\frac{\partial\rho(\vec{r},t)}{\partial t}+\vec{\nabla}\vec{\jmath}(\vec{r},t)=0, \tag{7}$$

where the current density takes a form of

$$\jmath^\alpha(\vec{r},t)=\sum_S\int dR\sum_j^N\delta(\vec{r}-\vec{r}_j)\frac{1}{2m_j}(\hat{D}_j^{+\alpha}\psi_s^+(R,t)\psi_s(R,t) \tag{8}$$

$$+\psi_s^+(R,t)\hat{D}_j^\alpha\psi_s(R,t)),$$

A momentum balance equation can be derived by differentiating current density 8 with respect to time

$$\partial_t\jmath^\alpha(\vec{r},t)+\partial_\beta\Re^{\alpha\beta}(\vec{r},t)=q\rho(\vec{r},t)E_{ext}^\alpha(\vec{r},t) \tag{9}$$

$$+\frac{q}{mc}\varepsilon^{\alpha\beta\gamma}\jmath_\beta(\vec{r},t)B_{ext}^\gamma(\vec{r},t)-\int d\vec{r}'\,q^2\partial^\alpha G(\vec{r},\vec{r}')\rho_2(\vec{r},\vec{r}',t)$$

$$+M_\beta(\vec{r},t)\partial^\alpha B_{ext}^\beta(\vec{r},t)+\int d\vec{r}'\,\partial^\alpha F^{\gamma\delta}(\vec{r},\vec{r}')M^{\gamma\delta}(\vec{r},\vec{r}',t).$$

$$\Re^{\alpha\beta}(\vec{r},t)=\sum_S\int dR\sum_{j=1}^N\delta(\vec{r}-\vec{r}_j)\frac{1}{4m_j}(\psi^+(R,t)\hat{D}_j^\alpha D_j^\beta\psi(R,t) \tag{10}$$

$$+(\hat{D}_j^\alpha\psi(R,t))^+D_j^\beta\psi(R,t)+h.c.)$$

represents the momentum current density tensor.

Momentum balance equation 9 contains the particle magnetic moment density[1]

$$M^\alpha(\vec{r},t)=\sum_s\int dR\sum_{j=1}^N\delta(\vec{r}-\vec{r}_j)\psi_s^+\hat{\sigma}_j^\alpha\psi_s, \tag{11}$$

The *Coulomb* and $Spin-Spin$ interactions between the particles are represented in Eq. 9 by the terms where

$$\rho_2(\vec{r},\vec{r}',t)=\sum_s\int dR\sum_{j\neq k}^N\delta(\vec{r}-\vec{r}_j)\delta(\vec{r}'-\vec{r}_k)\psi^+(R,t)\psi(R,t), \tag{12}$$

is the two-particle probability density for the occurrence of two particle in the neighborhoods of the points $\vec{r}$ and $\vec{r}^{\,\prime}$ normalized by $N(N-1)$, and two-particle tensor of the magnetic moment density

$$M^{\alpha\beta}=\sum_{s}\int dR\sum_{j\neq k}^{N}\delta(\vec{r}-\vec{r}_j)\delta(\vec{r}^{\,\prime}-\vec{r}_k)\mu_j\mu_k\psi^{+}\hat{\sigma}_j^{\alpha}\hat{\sigma}_k^{\beta}\psi(R,t). \tag{13}$$

Differentiation of M^{α} with respect to time and applying of the Schrodinger equation with Hamiltonian 3 leads to magnetization equation. The equation representing the non-relativistic evolution of $spin-1/2$ motion takes a form of

$$\partial_t M^{\alpha}(\vec{r},t)+\partial_{\beta}\Im_M^{\alpha\beta}=\frac{2\mu}{\hbar}\epsilon^{\alpha\beta\gamma}M^{\beta}(\vec{r},t)B_{ext}^{\gamma}(\vec{r},t)$$

$$+\frac{2\mu}{\hbar}\epsilon^{\alpha\beta\gamma}\int d\vec{r}^{\,\prime}\partial^{\alpha}F^{\gamma\delta}(\vec{r},\vec{r}^{\,\prime})M^{\beta\delta}(\vec{r},\vec{r}^{\,\prime},t)$$

where the tensor of the magnetic moment flux density is

$$\Im_M^{\alpha\beta}(\vec{r},t)=\sum_{S}\int dR\sum_{j=1}^{N}\delta(\vec{r}-\vec{r}_j)\frac{\mu_j}{4m_j}(\psi^{+}\hat{\sigma}_j^{\alpha}\hat{D}_j^{\beta}\psi+ \tag{14}$$

$$+(\hat{\sigma}_j^{\alpha}D_j^{\beta}\psi)^{+}\psi)(R,t).$$

A. The vorticity and spin vorticity equation

The spinning quantum magnetohydrodynamics should be explain the vorticity evolution. The main idea of this paper was to create the hydrodynamics foundation for the vortex dynamic in the context of spinning quantum plasma. We use the MPQHD approach to receive the equations for the particle Vorticity density obeying the standard vortex dynamics. We determine the vorticity density vector of particles in the neighborhood of $\vec{r}$ in a physical space as

$$\Omega^{\alpha}(\vec{r},t)=\sum_{S}\int dR\sum_{j}^{N}\delta(\vec{r}-\vec{r}_j)\frac{\varepsilon^{\alpha\beta\gamma}}{2m_j}\hat{\nabla}_{j\beta}(\hat{D}_j^{+\gamma}\psi_s^{+}\psi_s \tag{15}$$

$$+\psi_s^{+}\hat{D}_j^{\gamma}\psi_s)(R,t),$$

The derivation of the quantum vorticity equation that include the spin is similar to the previous section

$$\partial_t\Omega^{\alpha}(\vec{r},t)+\varepsilon^{\alpha\beta\gamma}\partial_{\beta}\partial_k\Re^{k\gamma}(\vec{r},t)=F^{\alpha}(\vec{r},t). \tag{16}$$

Lets introduce a non trivial the spin or quantum vorticity as

$$\Theta_S^{\alpha}(\vec{r},t)=\sum_{S}\int dR\sum_{j}^{N}\delta(\vec{r}-\vec{r}_j)\varepsilon^{\alpha\beta\gamma}\hat{\nabla}_{j\beta}(\psi_s^{+}\hat{\sigma}_{\gamma}\psi_s), \tag{17}$$

The equation for spin vorticity can be easily derived using the previous procedures

$$\partial_t\Theta_S^{\alpha}(\vec{r},t)+\varepsilon^{\alpha\beta\gamma}\partial_{\beta}\partial_k\Im^{k\gamma}(\vec{r},t)=\aleph^{\alpha}(\vec{r},t). \tag{18}$$

B. Velocity field

The velocity of $j-th$ particle $\vec{v}_j$ is determined by equation

$$\vec{v}_j=\frac{1}{m_j}(\vec{\nabla}_j S-i\hbar\varphi^{+}\vec{\nabla}_j\varphi)-\frac{q_j}{m_j c}\vec{A}_j, \tag{19}$$

The quantity $\vec{v}_j(R,t)$ describe the current of probability connected with the motion of $j-th$ particle, in general case $\vec{v}_j(R,t)$ depend on coordinate of all particles of the system R, where R is the totality of $3N$ coordinate of N particles of the system $R=(\vec{r}_1,...,\vec{r}_N)$.

The $S(R,t)$ value in the formula 19 represents the phase of the wave function and as the electron has spin, the wave function is now be expressed in the form .

$$\psi(R,t)=a(R,t)e^{\frac{iS}{\hbar}}\varphi(\vec{r},R,t), \tag{20}$$

where φ, normalized such that $\varphi^{+}\varphi=1$, is the new spinor, defined in the local frame of reference with the origin at the point $\vec{r}$. The spinor gives the spin part of the wave function.

We substituted the wave function in the definition of the basic hydrodynamical quantities. The momentum current density tensor 10 has the new form of

$$\Re^{\alpha\beta}(\vec{r},t)=m\rho(\vec{r},t)\upsilon^{\alpha}(\vec{r},t)\upsilon^{\beta}(\vec{r},t)+\wp^{\alpha\beta}(\vec{r},t)+ \tag{21}$$

$$+\Lambda^{\alpha\beta}(\vec{r},t)+\Upsilon_s^{\alpha\beta}(\vec{r},t),$$

where

$$\wp^{\alpha\beta}(\vec{r},t)=\sum_{S}\int dR\sum_{j=1}^{N}\delta(\vec{r}-\vec{r}_j)a^2m_ju_j^{\alpha}u_j^{\beta}, \tag{22}$$

is the well known kinetic pressure tensor. Value $u_j^{\alpha}(\vec{r},R,t)$ is a quantum equivalent of the thermal speed and $u_j^{\alpha}(\vec{r},R,t)=\upsilon_j^{\alpha}(R,t)-\upsilon(\vec{r},t)$.

The tensor $\Lambda^{\alpha\beta}$ is proportional to $\hbar^2$, has a purely quantum origin and can therefore be interpreted as an additional quantum pressure

$$\Lambda^{\alpha\beta}(\vec{r},t)=-\sum_{S}\int dR\sum_{j=1}^{N}\delta(\vec{r}-\vec{r}_j)a^2\frac{\hbar^2}{2m_j}\frac{\partial^2 lna}{\partial x_j^\alpha \partial x_j^\beta} \tag{23}$$

For the large system of noninteracting particles, this tensor is

$$\Lambda^{\alpha\beta}(\vec{r},t)=-\frac{\hbar^2}{4m}(\partial^\alpha\partial^\beta\rho(\vec{r},t)- \tag{24}$$

$$-\frac{1}{\rho(\vec{r},t)}\{\partial^\alpha\rho(\vec{r},t)\}\{\partial^\beta\rho(\vec{r},t)\})$$

The tensor $\Upsilon_s^{\alpha\beta}$ appears in the theory as a result of representations rotating electrons as an assembly of bodies continuously distributed in space. In the context of quantum hydrodynamics the force due to a new spin stress inside the fluid takes the form of

$$\Upsilon_s^{\alpha\beta}=-\frac{\hbar^2}{4m\mu^2}M_\gamma\partial^\alpha\partial^\beta(\frac{M^\gamma}{\rho}) \tag{25}$$

This new force emerges from the inhomogeneity of spin distribution and must be considered in the equation of motion, being the order of $\hbar^2$.

On the other hand, after the presentation of the wave function in the exponential form the tensor of the magnetic moment flux density takes the form of

$$\Im_M^{\alpha\beta}(\vec{r},t)=M^\alpha\upsilon^\beta(\vec{r},t)+\gamma_s^{\alpha\beta}(\vec{r},t), \tag{26}$$

where

$$\gamma_s^{\alpha\beta}(\vec{r},t)=\sum_{S}\int dR\sum_{j=1}^{N}\delta(\vec{r}-\vec{r}_j)\frac{\mu_j}{4m_j}(\varphi^+\hat{\sigma}_j^\alpha\hat{D}_j^\beta\varphi+ \tag{27}$$

$$+(\hat{\sigma}_j^\alpha D_j^\beta\varphi)^+\varphi)(R,t)$$

differ from the $\Im_M^{\alpha\beta}$ only by the change of the spinor ψ in 14 to the spin part of the wave function. In the context of quantum hydrodynamics we have the additional spin torque

$$\gamma_s^{\alpha\beta}(\vec{r},t)=\frac{\hbar}{2m\mu}\varepsilon^{\alpha\gamma\lambda}M_\gamma\partial_\beta(\frac{M^\lambda}{\rho}), \tag{28}$$

Taken in the approximation of self-consistent field, from 9, 14 and 18 we have the set of MQHD equation for the *electrons* and *positrons*, continuity equation, momentum balance equation, magnetic moment density equation take the form

$$\partial_t\rho_p+\vec{\nabla}(\rho_p\vec{\upsilon}_p)=0, \tag{29}$$

$$m_p\rho_p(\partial_t+\upsilon^\beta\partial_\beta)\vec{\upsilon}_p=q_p\rho_p\vec{E}_{ext}+\frac{1}{c}\vec{j}_{pe}\times\vec{B}_{ext}-\vec{\nabla}\wp_p+\frac{\hbar^2}{2m^2}\rho_p\vec{\nabla}(\frac{\triangle\sqrt{\rho_p}}{\sqrt{\rho_p}})+M_{p\beta}\vec{\nabla}B_{ext}^\beta+\frac{\hbar^2}{4m\mu^2}\partial_\beta\{M_p^\gamma\vec{\nabla}\partial^\beta(\frac{M_p^\gamma}{\rho_p})\} \tag{30}$$

$$-\rho_p\vec{\nabla}\int d\vec{r}'q_p^2G(\vec{r},\vec{r}')\rho_p(\vec{r}',t)+M_{p\gamma}\vec{\nabla}\int d\vec{r}'F^{\gamma\delta}(\vec{r},\vec{r}')M_p^\delta(\vec{r}',t),$$

$$(\partial_t+\upsilon^\beta\partial_\beta)\vec{M}_p=\frac{2\mu_p}{\hbar}\vec{M}_p\times\vec{B}_{ext}+\frac{\hbar}{2m\mu_p}\partial_k\{\vec{M}_p\times\partial^k(\frac{\vec{M}_p}{\rho_p})\}+\frac{2\mu_p}{\hbar}\epsilon^{\alpha\beta\gamma}M_p^\beta\int d\vec{r}'F^{\gamma\delta}(\vec{r},\vec{r}')M_p^\delta(\vec{r}',t), \tag{31}$$

Lets discuss the physical significance of terms on the right side of the system of MQHD equations obtained above 29 - 32. The first and second terms in Eq. 30 describe the well known interaction with external electromagnetic field, where the first term represents the effect of external electric field on the charge density and the second term is the Lorenz force field. The fourth term is a quantum force produced by density fluctuations, which has its origin in the socalled Madelung potential. The fifth term appears in the equation of motion 30 through the magnetization energy and depends on the spin or magnetic moment density of particles. The sixth term represents the self-force or *magnetic moment density stress* inside the electron or positron fluid. This spin self-force appears even in absence of the electromagnetic fields and arises from the inhomogeneity of magnetic moment density distribution. Other terms in 30 describe a force field that represents interactions between particles, namely the *Coulomb* interaction of charges and $Spin-Spin$ interactions.

The second term in the equation of magnetic moment density motion 31 represents the effect additional *magnetic moment density torque* on the magnetic moment density evolution and tends to align spins parallel.

Manipulations of the Eq. 18 and momentum balance dynamical equation 30 yields the hydrodynamics vorticity density dynamical equation $\vec{\omega}_p = \vec{\Omega}_p/\rho_p$

$$\partial_t\vec{\omega}_p = \vec{\nabla}\times(\vec{v}_p\times\vec{\omega}_p) - \vec{\nabla}(\frac{1}{\rho_p})\times\vec{\nabla}\wp_p + \vec{\nabla}(\frac{M_{pk}}{\rho_p})\times\vec{\nabla}B^k_{ext} + \frac{1}{cm_p}\vec{\nabla}\times(\frac{1}{\rho_p}\vec{j}_{pe}\times\vec{B}_{ext}) \tag{32}$$

$$+\frac{\hbar^2}{4m_p^2\mu_p^2}\vec{\nabla}(\frac{M^\nu_p}{\rho_p})\times\vec{\nabla}\{\frac{1}{\rho_p}\nabla_k(\rho_p\nabla^k(\frac{M^\nu_p}{\rho_p}))\} + \vec{\nabla}(\frac{M_{p\gamma}}{\rho_p})\times\vec{\nabla}\int d\vec{r}^{\prime}F^{\gamma\delta}(\vec{r},\vec{r}^{\prime})M^\delta_p(\vec{r}^{\prime},t).$$

and manipulations of the Eq. 33 and magnetic moment density dynamical equation 31 yields the hydrodynamics quantum spin vorticity evolution equation $\vec{\varpi}_p = \vec{\Theta}_{Sp}/\rho_p$

$$\partial_t\vec{\varpi}_p = \vec{\nabla}\times(\vec{v}_p\times\vec{\varpi}_p) + \frac{2\mu_p}{\hbar}\vec{\nabla}\times(\frac{\vec{M}_p}{\rho_p}\times\vec{B}_{ext}) + \epsilon^{\alpha\beta\gamma}\varepsilon_{\gamma\mu\nu}\frac{2\mu_p}{\hbar}\nabla_\beta\{\frac{M^\mu_p}{\rho_p}\int d\vec{r}^{\prime}F^{\nu\delta}(\vec{r},\vec{r}^{\prime})M^\delta_p(\vec{r}^{\prime},t)\} \tag{33}$$

$$+\frac{\hbar}{2m\mu_p}\vec{\nabla}\times(\frac{\vec{M}_p}{\rho_p}\times\frac{1}{\rho_p}\nabla_k(\rho_p\nabla^k(\frac{M^\nu_p}{\rho_p}))$$

The vorticity density evolution equation 32 shows the different physical factors associated with the generation of vorticity. The second term on the right side of 32 is proportional to the gas pressure and is responsible for the hydrodynamic baroclinic vorticity generation the classical vortex field. The third term represents the magnetic baroclinic vorticity. The fourth term contains the information about the vorticity generated by the magnetic tension. The fifth term associated with the magnetic vorticity generation even in absence of the magnetic field. The seventh term characterizes the effect of the $Spin - Spin$ interactions on the vorticity evolution. Equation 32 contains the normal electron or positron current density $\vec{j}_{ep} = q_p\rho_p\vec{v}_p$, and the magnetic moment density $\vec{M}_p = \rho_p\vec{\mu}_p$.

The equations 30, 31 and 32 allow for the introduction of an internal potential and internal magnetic field[8] and can be considered like classical effects.

II. CONCLUSIONS

In this paper we analyzed *vorticity* excitations caused by the magnetic moment density dynamics in systems of charged $1/2 - spin$ particles. MQHD equations are a consequence of MPSE in which particles interaction is directly taken into account. In our work we consider the *Coulomb* and $Spin - Spin$ interactions. The system of MQHD equations we constructed is comprised by equations of continuity, of the momentum balance, of the magnetic moment density evolution and of the vorticity density dynamics. In our studies of wave processed we used a self-consistent field approximation of the MQHD equations.

The equations we are interested in, determining the system dynamics, are the hydrodynamic equations for the spinning plasma. This equations have an additional quantum contribution proportional to $\hbar^2$ and spin corrections, additional *Magnetic Moment Stress* and *Magnetic Moment Torque* which have been derived in the absence of (thermal) fluctuation of the spin about the macroscopic average. But in such a situation (thermal) effects on the spin might be important. The main objective of this paper was to construct an appropriate a new generalized quantum vorticity equation for spin quantum plasmas that contains the magnetic, non-magnetic terms and the spin dependent forces being non potential. Important that the quantum Madelung potential do not contribute to the vorticity evolution.

Appendix A

The terms represented *Coulomb* and $Spin - Spin$ interactions in Eq. 30, 31 and 32 leads to the appearance of the self-consistent electric field E_{int} and the self-consistent magnetic field B_{spin}

$$\vec{\nabla}\vec{E}_{int} = 4\pi q\rho, \qquad \vec{\nabla}\times\vec{E}_{int} = 0, \tag{A1a}$$

$$\vec{\nabla}\vec{B}_{spin} = 0, \qquad \vec{\nabla}\times\vec{B}_{spin} = 4\pi\vec{\nabla}\times\vec{M}.$$

Two-particle functions 12 and 13 have the ground expressions

$$\rho_2(\vec{r},\vec{r}^{\prime},t) = \rho(\vec{r},t)\rho(\vec{r}^{\prime},t) + \varrho(\vec{r},\vec{r}^{\prime},t), \tag{A2}$$

$$M_2^\alpha(\vec{r},\vec{r}^{\,\prime},t)=M^\alpha(\vec{r},t)M^\alpha(\vec{r}^{\,\prime},t)+\chi^\alpha(\vec{r},\vec{r}^{\,\prime},t), \qquad \text{(A3)}$$

where $\varrho(\vec{r},\vec{r}^{\,\prime},t)$ and $\chi^\alpha(\vec{r},\vec{r}^{\,\prime},t)$ - are the the correlation functions.

[1] L. S. Kuzmenkov, S. G. Maksimov, V. V. Fedoseev, Theor. Math. Phys. **126**, 110 (2001)

[2] L. S. Kuz'menkov, S. G. Maksimov, V. V. Fedoseev, Theor. Math. Phys. **126**, 212 (2001)

[3] M. Marklund and G. Brodin, Phys. Rev. Lett. **98**, 025001 (2007)

[4] G. Brodin and M. Marklund, New J. Phys. **9**, 277 (2007)

[5] M. Bonitz et al., J. Phys. A: Math. Gen. **36**, 5921 (2003)

[6] P. K. Shukla, Phys. Lett. A **369**, 312 (2007)

[7] P. K. Shukla, Nature Phys. **5**, 92 (2009)

[8] T. Takabayasi, Prog. Theor. Phys. **14**, 283 (1955)

[9] T. Takabayasi, J. P. Vigier, Prog. Theor. Phys. **18**, 573 (1957)

[10] T. Takabayasi, Prog. Theor. Phys. **70**, 1 (1983)

[11] T. Takabayasi, Prog. Theor. Phys. Suppl. **4**, 2 (1957)

[12] E. Madelung, Z. Phys. **40**, 322 (1927)

[13] L. S. Kuzmenkov, S. G. Maksimov, Theor. Math. Phys. **118**, 227 (1999)

[14] Swadesh M. Mahajan, Felipe A. Asenjo, Phys. Rev. Lett. **107**, 195003 (2011)

[15] S. Shelyag, P. Keys, M. Mathioudakis, F.P. Keenan, e-print arXiv:1010.5604 (2010)

[16] T. Emonet, F. Moreno-Insertis, ApJ **492**, 804 (1998)

[17] Anirban Guha, Mona Rahmani, and Gregory A. Lawrence, Phys. Rev. E **87**, 013020 (2013)

[18] Wouter J. T. Bos (LMFA), Shinpei Futatani (PIIM), Sadruddin Benkadda (PIIM), Marie Farge (LMD), Kai Schneider (M2P2), Physics of Plasmas **15**, 072305 (2008)

[19] P. K. Shukla and B. Eliasson, Phys. Rev. Lett. **96**, 245001 (2006)

[20] Stefanie Braun, Felipe A. Asenjo, and Swadesh M. Mahajan, Phys. Rev. Lett. **109**, 175003 (2013)